§610.16

dried vaccines when the accompanying diluent contains a preservative; or to an Allergenic Product in 50 percent or more volume in volume (v/v) glycerin. An adjuvant shall not be introduced into a product unless there is satisfactory evidence that it does not affect adversely the safety or potency of the product. The amount of aluminum in the recommended individual dose of a biological product shall not exceed:

- (1) 0.85 milligrams if determined by assay;
- (2) 1.14 milligrams if determined by calculation on the basis of the amount of aluminum compound added: or
- (3) 1.25 milligrams determined by assay provided that data demonstrating that the amount of aluminum used is safe and necessary to produce the intended effect are submitted to and approved by the Director, Center for Biologics Evaluation and Research or the Director, Center for Drug Evaluation and Research (see mailing addresses in §600.2 of this chapter).
- (b) Extraneous protein; cell culture produced vaccines. Extraneous protein known to be capable of producing allergenic effects in human subjects shall not be added to a final virus medium of cell culture produced vaccines intended for injection. If serum is used at any stage, its calculated concentration in the final medium shall not exceed 1:1000 000
- (c) Antibiotics. A minimum concentration of antibiotics, other than penicillin, may be added to the production substrate of viral vaccines.
- (d) The Director of the Center for Biologics Evaluation and Research or the Director of the Center for Drug Evaluation and Research may approve an exception or alternative to any requirement in this section. Requests for such exceptions or alternatives must be in writing.

[38 FR 32056, Nov. 20, 1973, as amended at 46 FR 51903, Oct. 23, 1981; 48 FR 13025, Mar. 29, 1983; 48 FR 37023, Aug. 16, 1983; 49 FR 23834, June 8, 1984; 50 FR 4134, Jan. 29, 1985; 51 FR 15607, Apr. 25, 1986; 55 FR 11013, Mar. 26, 1990; 70 FR 14985, Mar. 24, 2005; 76 FR 20518, Apr. 13, 2011]

§610.16 Total solids in serums.

Except as otherwise provided by regulation, no liquid serum or antitoxin shall contain more than 20 percent total solids.

§ 610.17 Permissible combinations.

Licensed products may not be combined with other licensed products either therapeutic, prophylactic or diagnostic, except as a license is obtained for the combined product. Licensed products may not be combined with nonlicensable therapeutic, prophylactic, or diagnostic substances except as a license is obtained for such combination.

§610.18 Cultures.

- (a) Storage and maintenance. Cultures used in the manufacture of products shall be stored in a secure and orderly manner, at a temperature and by a method that will retain the initial characteristics of the organisms and insure freedom from contamination and deterioration.
- (b) Identity and verification. Each culture shall be clearly identified as to source strain. A complete identification of the strain shall be made for each new stock culture preparation. Primary and subsequent seed lots shall be identified by lot number and date of preparation. Periodic tests shall be performed as often as necessary to verify the integrity of the strain characteristics and freedom from extraneous organisms. Results of all periodic tests for verification of cultures and determination of freedom from extraneous organisms shall be recorded and retained.
- (c) Cell lines used for manufacturing biological products—(1) General requirements. Cell lines used for manufacturing biological products shall be:
 - (i) Identified by history;
- (ii) Described with respect to cytogenetic characteristics and tumorigenicity;
- (iii) Characterized with respect to in vitro growth characteristics and life potential; and
- (iv) Tested for the presence of detectable microbial agents.
- (2) Tests. Tests that are necessary to assure the safety, purity, and potency of a product may be required by the Director, Center for Biologics Evaluation